

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 September 2005 (15.09.2005)

PCT

(10) International Publication Number
WO 2005/084277 A2

(51) International Patent Classification: Not classified

(21) International Application Number:
PCT/US2005/006448

(22) International Filing Date: 28 February 2005 (28.02.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/548,769 27 February 2004 (27.02.2004) US

(71) Applicant (for all designated States except US): **UNIVERSITY OF DENVER** [US/US]; 2240 East Buechel Blvd.
4th Floor, Denver, CO 80208 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ST. JOHN, Judith**

[US/US]; 7090 W. Radcliff Avenue, Littleton, CO 80123 (US). **QUINN, Thomas, W.** [US/US]; 3515 W. 115th Avenue, Westminster, CO 80031 (US).


(74) Agents: **TRAVER, Robert, D.** et al.; Sheridan Ross P.C., 1560 Broadway, Suite 1200, Denver, CO 80202-5141 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,

[Continued on next page]

(54) Title: METHOD OF ISOLATING NUCLEIC ACID TARGETS



(57) Abstract: The invention provides efficient methods of isolating specific nucleic acid targets to obtain information from target nucleic acid sequences in a relatively short time period. DNA or cDNA is enzymatically digested into smaller fragments, double-stranded DNA linkers are added onto the ends of the DNA fragments to flank each fragment with a known DNA sequence. The fragments are mixed with an oligonucleotide probe that is bound to a marker and contains a conserved nucleic acid sequence of interest. The fragments that hybridize to the probe through nucleotide base pair complementation become indirectly connected to the marker. These target fragments are captured using a capture agent that specifically recognizes the marker and treated to prevent non-specific binding. Captured fragments are typically cloned prior to sequencing. The captured fragments may also be amplified using PCR to increase the efficiency of the cloning.



ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- *without international search report and to be republished upon receipt of that report*